

Implications of CHOOZ results for the decoupling of solar and atmospheric neutrino oscillations

S.M. Bilenky

*Joint Institute for Nuclear Research, Dubna, Russia, and
INFN, Sezione di Torino, and Dipartimento di Fisica Teorica, Università di Torino,
Via P. Giuria 1, I-10125 Torino, Italy*

C. Giunti

*INFN, Sezione di Torino, and Dipartimento di Fisica Teorica, Università di Torino,
Via P. Giuria 1, I-10125 Torino, Italy*

Abstract

We have considered the results of solar and atmospheric neutrino oscillation experiments in the scheme of mixing of three neutrinos with a mass hierarchy. It is shown that the recent results of the CHOOZ experiment imply that $|U_{e3}|^2 \ll 1$ (U is the neutrino mixing matrix), that the oscillations of solar neutrinos are described by the two-generation formalism and that the oscillations of solar and atmospheric neutrinos decouple. It is also shown that if not only $|U_{e3}|^2 \ll 1$ but also $|U_{e3}| \ll 1$, then the oscillations of atmospheric neutrinos do not depend on matter effects and are described by the two-generation formalism. In this case, with an appropriate identification of the mixing parameters, the two-generation analyses of solar and atmospheric neutrino data provide direct information on the mixing parameters of three neutrinos. We discuss the possibility to get information on $|U_{e3}|^2$ in long-baseline neutrino oscillation experiments.

PACS numbers: 14.60.Pq, 26.65.+t, 95.85.Ry

I. INTRODUCTION

The existence of the solar [1] and atmospheric [2] neutrino problems have received in the last months impressive confirmations from the preliminary results of the SuperKamiokande experiment [3,4]. The traditional analyses of solar [5–8,3] and atmospheric [9–11,4] neutrino data in terms of neutrino oscillations (see [12–14]) have been performed under the assumption of two-neutrino mixing. In particular, this assumption has been adopted in the most recent analyses of the solar [15,16] and atmospheric [17] neutrino data, which include preliminary data from SuperKamiokande [3,4]. However, we know that there are three light flavor neutrinos, ν_e , ν_μ , ν_τ , that can participate to the oscillations of solar and atmospheric neutrinos and it is reasonable to ask what information on the mixing of three neutrinos can be obtained from the two-generation analyses of solar and atmospheric data.

Here we consider a scheme of mixing of three massive neutrinos with the mass hierarchy

$$m_1 \ll m_2 \ll m_3, \quad (1)$$

which is motivated by the see-saw mechanism [18] and by the analogy with the mass spectra of charged leptons and up and down quarks. The masses m_k ($k = 1, 2, 3$) are associated to the massive neutrino fields ν_k , whose left-handed components ν_{kL} are connected with the left-handed components $\nu_{\alpha L}$ ($\alpha = e, \mu, \tau$) of the flavor neutrino fields by the relation

$$\nu_{\alpha L} = \sum_{k=1}^3 U_{\alpha k} \nu_{kL}, \quad (2)$$

where U is a 3×3 unitary mixing matrix. In this scheme there are two independent mass-squared differences, $\Delta m_{21}^2 \equiv m_2^2 - m_1^2$ and $\Delta m_{31}^2 \equiv m_3^2 - m_1^2$, that generate neutrino oscillations for two different scales of the ratio E/L , where E is the neutrino energy and L is the distance of propagation. The solar and atmospheric neutrino anomalies can be explained in terms of neutrino oscillations only if Δm_{21}^2 and Δm_{31}^2 are relevant, respectively, for the oscillations of solar and atmospheric neutrinos [19–24].

In Sections II and III we will show that in the scheme under consideration the recent results of the CHOOZ experiment [25] imply that $|U_{e3}|^2 \ll 1$, that the oscillations of solar neutrinos are described by the two-generation formalism and that the oscillations of solar and atmospheric neutrinos depend on different and independent elements of the neutrino mixing matrix, *i.e.* they decouple. In Section III it is shown that if not only $|U_{e3}|^2 \ll 1$ but also $|U_{e3}| \ll 1$, then the oscillations of atmospheric neutrinos do not depend on matter effects and are described by the two-generation formalism. In this case, with an appropriate identification of the mixing parameters, the formalism used in the two-generation analyses of solar and atmospheric neutrino data is appropriate also in the case of three-neutrino mixing and the results of these analyses provide direct information on the mixing parameters of three neutrinos. In Section IV we discuss the implications of the present solar and atmospheric neutrino data for the values of the elements of the neutrino mixing matrix. As shown in Section V, the hypothesis $|U_{e3}| \ll 1$ could be tested by future accelerator long-baseline neutrino oscillation experiments.

In this paper we do not consider the indication in favor of neutrino oscillations obtained by the LSND experiment [26], which needs the enlargement of the neutrino mixing scheme

with the introduction of a sterile neutrino (see [27,28]). A possible decoupling of solar and atmospheric neutrino oscillations in this enlarged scheme will be discussed elsewhere.

II. CHOOZ AND SOLAR NEUTRINOS

The first reactor long-baseline neutrino oscillation experiment CHOOZ found no evidence for neutrino oscillations [25]. The CHOOZ collaboration has published an exclusion curve in the plane of the two-generation mixing parameters which shows that

$$\sin^2 2\vartheta_{\text{CHOOZ}} \leq 0.18 \quad \text{for} \quad \Delta m_{\text{CHOOZ}}^2 \gtrsim 10^{-3} \text{ eV}^2. \quad (3)$$

In the three-neutrino scheme under consideration these parameters are given by (see [23,24])

$$\sin^2 2\vartheta_{\text{CHOOZ}} = 4|U_{e3}|^2(1 - |U_{e3}|^2) \quad \text{and} \quad \Delta m_{\text{CHOOZ}}^2 = \Delta m_{31}^2. \quad (4)$$

If $\Delta m_{31}^2 > 10^{-3} \text{ eV}^2$, as indicated by the solution of the atmospheric neutrino anomaly [9], the upper bound (4) on $\sin^2 2\vartheta_{\text{CHOOZ}}$ implies that

$$|U_{e3}|^2 \leq 5 \times 10^{-2} \quad \text{or} \quad |U_{e3}|^2 \geq 0.95. \quad (5)$$

A large value of $|U_{e3}|^2$ does not allow to explain the solar neutrino problem with neutrino oscillations (see [29]). Indeed, in the scheme under consideration the averaged survival probability of solar electron neutrinos is given by [19]

$$P_{\nu_e \rightarrow \nu_e}^{\text{sun}}(E) = (1 - |U_{e3}|^2)^2 P_{\nu_e \rightarrow \nu_e}^{(1,2)}(E) + |U_{e3}|^4, \quad (6)$$

where E is the neutrino energy and $P_{\nu_e \rightarrow \nu_e}^{(1,2)}(E)$ is the two-generation survival probability of solar ν_e 's which depends on

$$\Delta m_{\text{sun}}^2 = \Delta m_{21}^2 \quad \text{and} \quad \sin \vartheta_{\text{sun}} = \frac{|U_{e2}|}{\sqrt{1 - |U_{e3}|^2}}. \quad (7)$$

The expression (6) implies that $P_{\nu_e \rightarrow \nu_e}^{\text{sun}}(E) \geq |U_{e3}|^4$ and for $|U_{e3}|^2 \geq 0.95$ we have $P_{\nu_e \rightarrow \nu_e}^{\text{sun}}(E) \geq 0.90$. With such a high and practically constant value of $P_{\nu_e \rightarrow \nu_e}^{\text{sun}}(E)$ it is not possible to explain the suppression of the solar ν_e flux measured by all experiments (Homestake [5], Kamiokande [6], GALLEX [7], SAGE [8] and SuperKamiokande [3]) with respect to that predicted by the Standard Solar Model [30–32].

Therefore, we are led to the conclusion that the results of the CHOOZ experiment imply that $|U_{e3}|^2$ is small:

$$|U_{e3}|^2 \leq 5 \times 10^{-2}. \quad (8)$$

Furthermore, from Eqs.(6) and (7) one can see that for such small values of $|U_{e3}|^2$ we have

$$P_{\nu_e \rightarrow \nu_e}^{\text{sun}}(E) \simeq P_{\nu_e \rightarrow \nu_e}^{(1,2)}(E) \quad \text{and} \quad \sin \vartheta_{\text{sun}} \simeq |U_{e2}|. \quad (9)$$

Hence the two-generation analyses of the solar neutrino data are appropriate in the three-neutrino scheme with a mass hierarchy and they give information on the values of

$$\Delta m_{21}^2 = \Delta m_{\text{sun}}^2 \quad \text{and} \quad |U_{e2}| \simeq \sin \vartheta_{\text{sun}}. \quad (10)$$

According to the most recent analysis of the solar neutrino data [16], which include preliminary data from SuperKamiokande [3], the ranges of the mixing parameters allowed at 90% CL for the small and large mixing angle MSW [33] solutions and for the vacuum oscillation solution are, respectively,

$$4 \times 10^{-6} \text{ eV}^2 \lesssim \Delta m_{\text{sun}}^2 \lesssim 1.2 \times 10^{-5} \text{ eV}^2, \quad 3 \times 10^{-3} \lesssim \sin^2 2\vartheta_{\text{sun}} \lesssim 1.1 \times 10^{-2}, \quad (11)$$

$$8 \times 10^{-6} \text{ eV}^2 \lesssim \Delta m_{\text{sun}}^2 \lesssim 3.0 \times 10^{-5} \text{ eV}^2, \quad 0.42 \lesssim \sin^2 2\vartheta_{\text{sun}} \lesssim 0.74, \quad (12)$$

$$6 \times 10^{-11} \text{ eV}^2 \lesssim \Delta m_{\text{sun}}^2 \lesssim 1.1 \times 10^{-10} \text{ eV}^2, \quad 0.70 \lesssim \sin^2 2\vartheta_{\text{sun}} \leq 1. \quad (13)$$

Therefore, taking into account that $|U_{e3}|^2$ is small and assuming that $|U_{e2}| \leq |U_{e1}|$ (this choice is necessary only for the MSW solutions), we have

$$|U_{e1}| \simeq 1, \quad |U_{e2}| \simeq 0.03 - 0.05 \quad (\text{small mixing MSW}), \quad (14)$$

$$|U_{e1}| \simeq 0.87 - 0.94, \quad |U_{e2}| \simeq 0.35 - 0.49 \quad (\text{large mixing MSW}), \quad (15)$$

$$|U_{e1}| \simeq 0.71 - 0.88, \quad |U_{e2}| \simeq 0.48 - 0.71 \quad (\text{vacuum oscillations}). \quad (16)$$

Notice that these ranges are statistically rather stable. For example, the range of $\sin^2 2\vartheta_{\text{sun}}$ allowed at 99% CL in the case of the large mixing angle MSW solution is $0.36 \lesssim \sin^2 2\vartheta_{\text{sun}} \lesssim 0.85$ [16], which imply $|U_{e1}| \simeq 0.83 - 0.95$, $|U_{e2}| \simeq 0.32 - 0.55$ (confront with Eq.(15)).

III. ATMOSPHERIC NEUTRINOS

The evolution equation for the flavor amplitudes ψ_α ($\alpha = e, \mu, \tau$) of atmospheric neutrinos propagating in the interior of the Earth can be written as (see [34,24])

$$i \frac{d}{dt} \Psi = \frac{1}{2E} (U M^2 U^\dagger + A) \Psi, \quad (17)$$

with

$$\Psi \equiv \begin{pmatrix} \psi_e \\ \psi_\mu \\ \psi_\tau \end{pmatrix}, \quad M^2 \equiv \text{diag}(0, \Delta m_{21}^2, \Delta m_{31}^2), \quad A \equiv \text{diag}(A_{CC}, 0, 0), \quad (18)$$

and $A_{CC} \equiv 2EV_{CC}$, where $V_{CC} = \sqrt{2}G_F N_e$ is the charged-current effective potential which depends on the electron number density N_e of the medium (G_F is the Fermi constant and for anti-neutrinos A_{CC} must be replaced by $\bar{A}_{CC} = -A_{CC}$). If the squared-mass difference Δm_{21}^2 is relevant for the explanation of the solar neutrino problem, we have

$$\frac{\Delta m_{21}^2 R_\oplus}{2E} \ll 1, \quad (19)$$

where $R_{\oplus} = 6371 \text{ Km}$ is the radius of the Earth. Notice, however, that caution is needed for low-energy atmospheric neutrinos if $\Delta m_{21}^2 \gtrsim 10^{-5} \text{ eV}^2$, as in the case of the large mixing angle MSW solution of the solar neutrino problem and marginally in the case of the small mixing angle MSW solution (see Eqs.(12) and (11)). Indeed, if $\Delta m_{21}^2 \gtrsim 10^{-5} \text{ eV}^2$ we have $\Delta m_{21}^2 R_{\oplus}/2E \ll 1$ only for $E \gg 150 \text{ MeV}$. In this case, in order to get information on the three-neutrino mixing matrix with a two-generation analysis it is necessary to analyze the atmospheric neutrino data with a cut in energy such that $\Delta m_{21}^2 R_{\oplus}/2E \ll 1$. In order to be on the safe side, when we will consider the case of the MSW solutions of the solar neutrino problem we will take into account the information obtained from the two-generation fit of the preliminary SuperKamiokande multi-GeV data alone (Figure 11 of [17]).

The inequalities (19) imply that the phase generated by Δm_{21}^2 can be neglected for atmospheric neutrinos and M^2 can be approximated with

$$M^2 \simeq \text{diag}(0, 0, \Delta m_{31}^2). \quad (20)$$

In this case (taking into account that the phases of the matrix elements $U_{\alpha 3}$ can be included in the charged lepton fields) we have

$$(U M^2 U^\dagger)_{\alpha'\alpha} \simeq \Delta m_{31}^2 |U_{\alpha'3}| |U_{\alpha 3}|. \quad (21)$$

Comparing this expression with Eqs.(9) and (10), one can see that the oscillations of solar and atmospheric neutrinos depend on different and independent Δm^2 's and on different and independent elements of the mixing matrix, *i.e.* they are decoupled. Strictly speaking $|U_{e2}|$ in Eqs.(9) and (10) is not independent from $|U_{e3}|$ because of the unitarity constraint $|U_{e1}|^2 + |U_{e2}|^2 + |U_{e3}|^2 = 1$, but the limit (8) on $|U_{e3}|^2$ implies that its contribution to the unitarity constraint is negligible.

Hence, we have shown that the smallness of $|U_{e3}|^2$ inferred from the results of the CHOOZ experiment imply that the oscillations of solar and atmospheric neutrinos are decoupled.

From Eqs.(17) and (21) one can see that unless $|U_{e3}| \ll 1$, the evolution equations of the atmospheric electron neutrino amplitude ψ_e and those of the muon and tau neutrino amplitudes ψ_μ and ψ_τ are coupled. In this case matter effects can contribute to the dominant $\nu_\mu \rightarrow \nu_\tau$ oscillations (see [35]) and the atmospheric neutrino data must be analyzed with the three generation evolution equation (17).

From the results of the CHOOZ experiment it follows that the quantity $|U_{e3}|^2$ is small and satisfy the inequality (8) (for $\Delta m_{31}^2 \gtrsim 10^{-3} \text{ eV}^2$). However, the upper bound for $|U_{e3}|$ implied by Eq.(8) is not very strong: $|U_{e3}| < 0.22$. In the following we will assume that not only $|U_{e3}|^2 \ll 1$, but also the element $|U_{e3}|$ that connects the first and third generations is small: $|U_{e3}| \ll 1$ (let us remind that in the quark sector $2 \times 10^{-2} \leq |V_{ub}| \leq 5 \times 10^{-2}$). We will consider the other elements of the mixing matrix as free parameters and we will see that these parameters can be determined by the two-neutrino analyses of the solar and atmospheric neutrino data. In Section V it will be shown that the hypothesis $|U_{e3}| \ll 1$ can be tested in future long-baseline neutrino oscillation experiments.

If $|U_{e3}| \ll 1$, for the evolution operator in Eq.(17) we have the approximate expression

$$U M^2 U^\dagger + A \simeq \Delta m_{31}^2 \begin{pmatrix} \frac{A_{CC}}{\Delta m_{31}^2} & 0 & 0 \\ 0 & |U_{\mu 3}|^2 & |U_{\mu 3}| |U_{\tau 3}| \\ 0 & |U_{\tau 3}| |U_{\mu 3}| & |U_{\tau 3}|^2 \end{pmatrix} \quad (22)$$

which shows that the evolution of ν_e is decoupled from the evolution of ν_μ and ν_τ . Thus, the survival probability of atmospheric ν_e 's is equal to one and $\nu_\mu \rightarrow \nu_\tau$ transitions are independent from matter effects and are described by a two-generation formalism. In this case, the two-generation analyses of the atmospheric neutrino data in terms of $\nu_\mu \rightarrow \nu_\tau$ are appropriate in the three-neutrino scheme under consideration and yield information on the values of the parameters

$$\Delta m_{31}^2 = \Delta m_{\text{atm}}^2 \quad \text{and} \quad |U_{\mu 3}| = \sin \vartheta_{\text{atm}}. \quad (23)$$

According to a recent analysis [17] of the atmospheric neutrino data, the ranges of Δm_{atm}^2 and $\sin^2 2\vartheta_{\text{atm}}$ for $\nu_\mu \rightarrow \nu_\tau$ oscillations allowed at 90% CL by the SuperKamiokande multi-GeV data and by all data are, respectively,

$$4 \times 10^{-4} \text{ eV}^2 \lesssim \Delta m_{\text{atm}}^2 \lesssim 8 \times 10^{-3} \text{ eV}^2, \quad 0.72 \lesssim \sin^2 2\vartheta_{\text{atm}} \leq 1, \quad (24)$$

$$4 \times 10^{-4} \text{ eV}^2 \lesssim \Delta m_{\text{atm}}^2 \lesssim 6 \times 10^{-3} \text{ eV}^2, \quad 0.76 \lesssim \sin^2 2\vartheta_{\text{atm}} \leq 1. \quad (25)$$

Thus, assuming that $|U_{\mu 3}| \leq |U_{\tau 3}|$ and taking into account the comments after Eq.(19), we have

$$|U_{\mu 3}| \simeq 0.49 - 0.71, \quad |U_{\tau 3}| \simeq 0.71 - 0.87 \quad (\text{MSW}), \quad (26)$$

$$|U_{\mu 3}| \simeq 0.51 - 0.71, \quad |U_{\tau 3}| \simeq 0.71 - 0.86 \quad (\text{vacuum oscillations}). \quad (27)$$

As in the case of the ranges (14)–(16), also the ranges (26)–(27) are statistically rather stable. For example, the range of $\sin^2 2\vartheta_{\text{sun}}$ allowed at 99% CL by all the atmospheric neutrino data is $0.66 \lesssim \sin^2 2\vartheta_{\text{sun}} \leq 1$ [17], which imply $|U_{\mu 3}| \simeq 0.46 - 0.71$, $|U_\tau| \simeq 0.71 - 0.89$ (confront with Eq.(27)).

IV. THE MIXING MATRIX

Taking into account the unitarity of the mixing matrix, the information in Eqs.(14)–(16) and (26)–(27), together with the assumption $|U_{e3}| \ll 1$, allow to infer the allowed ranges for the values of $|U_{\mu 1}|$, $|U_{\mu 2}|$, $|U_{\tau 1}|$ and $|U_{\tau 2}|$. The simplest way to do it is to start from the Maiani parameterization of a 3×3 mixing matrix [36]:

$$U = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13} & c_{12}c_{23} - s_{12}s_{23}s_{13} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13} & -c_{12}s_{23} - s_{12}c_{23}s_{13} & c_{23}c_{13} \end{pmatrix}, \quad (28)$$

where $c_{ij} \equiv \cos \vartheta_{ij}$ and $s_{ij} \equiv \sin \vartheta_{ij}$ and we have omitted possible CP-violating phases (one for Dirac neutrinos and three for Majorana neutrinos) on which there is no information.

A very small $|U_{e3}|$ implies that $|s_{13}| \ll 1$. In this case we have

$$U \simeq \begin{pmatrix} c_{12} & s_{12} & \ll 1 \\ -s_{12}c_{23} & c_{12}c_{23} & s_{23} \\ s_{12}s_{23} & -c_{12}s_{23} & c_{23} \end{pmatrix}. \quad (29)$$

Using the information on $|s_{12}| \simeq |U_{e2}|$ and $|s_{23}| \simeq |U_{\mu 3}|$ given by Eqs.(14)–(16) and (26)–(27), for the moduli of the elements of the mixing matrix we obtain:

$$\text{Small mixing MSW: } \begin{pmatrix} \simeq 1 & 0.03 - 0.05 & \ll 1 \\ 0.02 - 0.05 & 0.71 - 0.87 & 0.49 - 0.71 \\ 0.01 - 0.04 & 0.48 - 0.71 & 0.71 - 0.87 \end{pmatrix}, \quad (30)$$

$$\text{Large mixing MSW: } \begin{pmatrix} 0.87 - 0.94 & 0.35 - 0.49 & \ll 1 \\ 0.25 - 0.43 & 0.61 - 0.82 & 0.49 - 0.71 \\ 0.17 - 0.35 & 0.42 - 0.66 & 0.71 - 0.87 \end{pmatrix}, \quad (31)$$

$$\text{Vacuum oscillations: } \begin{pmatrix} 0.71 - 0.88 & 0.48 - 0.71 & \ll 1 \\ 0.34 - 0.61 & 0.50 - 0.76 & 0.51 - 0.71 \\ 0.24 - 0.50 & 0.36 - 0.62 & 0.71 - 0.86 \end{pmatrix}. \quad (32)$$

Let us remark that in the case of the small mixing angle MSW solution of the solar neutrino problem $|U_{e3}| \ll 1$ could be of the same order of magnitude as $|U_{e2}|$.

It is interesting to notice that, because of the large mixing of ν_μ and ν_τ with ν_2 , the transitions of solar ν_e 's in ν_μ 's and ν_τ 's are of comparable magnitude. However, this phenomenon and the values of the entries in the (ν_μ, ν_τ) – (ν_1, ν_2) sector of the mixing matrix cannot be checked with solar neutrino experiments because the low-energy ν_μ 's and ν_τ 's coming from the sun can be detected only with neutral-current interactions, which are flavor-blind. Moreover, it will be very difficult to check the values of $|U_{\mu 1}|$, $|U_{\mu 2}|$, $|U_{\tau 1}|$ and $|U_{\tau 2}|$ in laboratory experiments because of the smallness of m_2 .

In the derivation of Eqs.(30)–(32) we have assumed that $|U_{e2}| \leq |U_{e1}|$ and $|U_{\mu 3}| \leq |U_{\tau 3}|$. The other possibilities, $|U_{e2}| \geq |U_{e1}|$ and $|U_{\mu 3}| \geq |U_{\tau 3}|$, are equivalent, respectively, to an exchange of the first and second columns and to an exchange of the second and third rows in the matrices (30)–(32). Unfortunately, these alternatives are hard to distinguish experimentally because of the above mentioned difficulty to measure directly the values of $|U_{\mu 1}|$, $|U_{\mu 2}|$, $|U_{\tau 1}|$ and $|U_{\tau 2}|$. Only the choice $|U_{e2}| \leq |U_{e1}|$, which is necessary for the MSW solutions of the solar neutrino problem, could be confirmed by the results of the new generation of solar neutrino experiments (SuperKamiokande, SNO, ICARUS, Borexino, GNO and others [37]) if they will allow to exclude the vacuum oscillation solution.

V. ACCELERATOR LONG-BASELINE EXPERIMENTS

Future results from reactor long-baseline neutrino oscillation experiments (CHOOZ [25], Palo Verde [38], Kam-Land [39]) could allow to improve the upper bound (8) on $|U_{e3}|^2$. In this section we discuss how an improvement of this upper bound could be obtained by future accelerator long-baseline neutrino oscillation experiments that are sensitive to $\nu_\mu \rightarrow \nu_e$ transitions (K2K [40], MINOS [41], ICARUS [42] and others [43]).

If matter effects are not important, in the scheme under consideration the parameter $\sin^2 2\vartheta_{\mu e}$ measured in $\nu_\mu \rightarrow \nu_e$ long-baseline experiments is given by (see [23,24])

$$\sin^2 2\vartheta_{\mu e} = 4|U_{e3}|^2|U_{\mu 3}|^2. \quad (33)$$

If accelerator long-baseline neutrino oscillation experiments will not observe $\nu_\mu \rightarrow \nu_e$ transitions and will place an upper bound $\sin^2 2\vartheta_{\mu e} \leq \sin^2 2\vartheta_{\mu e}^{(\max)}$, it will be possible to obtain the limit

$$|U_{e3}|^2 \leq \frac{\sin^2 2\vartheta_{\mu e}^{(\max)}}{4|U_{\mu 3}|_{(\min)}^2}, \quad (34)$$

where $|U_{\mu 3}|_{(\min)}^2$ is the minimum value of $|U_{\mu 3}|^2$ allowed by the solution of the atmospheric neutrino anomaly and by the observation of $\nu_\mu \rightarrow \nu_\tau$ long-baseline transitions. For example, taking $|U_{\mu 3}|_{(\min)}^2 = 0.25$ (see Eq.(27)) we have $|U_{e3}|^2 \leq \sin^2 2\vartheta_{\mu e}^{(\max)}$. If a value of $\sin^2 2\vartheta_{\mu e}^{(\max)} \simeq 10^{-3}$, that corresponds to the sensitivity of the ICARUS experiment for one year of running [42], will be reached, it will be possible to put the upper bound $|U_{e3}| \lesssim 3 \times 10^{-2}$.

The observation of $\nu_\mu \rightarrow \nu_\tau$ transitions in long-baseline experiments will allow to establish a lower bound for $|U_{\mu 3}|^2$ because the parameter $\sin^2 2\vartheta_{\mu\tau}$ is given in the scheme under consideration by (see [23,24])

$$\sin^2 2\vartheta_{\mu\tau} = 4|U_{\mu 3}|^2 |U_{\tau 3}|^2. \quad (35)$$

From the unitarity relation $|U_{e3}|^2 + |U_{\mu 3}|^2 + |U_{\tau 3}|^2 = 1$ it follows that an experimental lower bound $\sin^2 2\vartheta_{\mu\tau} \geq \sin^2 2\vartheta_{\mu\tau}^{(\min)}$ allows to constraint $|U_{\mu 3}|^2$ in the range

$$\frac{1}{2} \left(1 - \sqrt{1 - \sin^2 2\vartheta_{\mu\tau}^{(\min)}} \right) \leq |U_{\mu 3}|^2 \leq \frac{1}{2} \left(1 + \sqrt{1 - \sin^2 2\vartheta_{\mu\tau}^{(\min)}} \right). \quad (36)$$

If $\sin^2 2\vartheta_{\mu\tau}^{(\min)}$ is found to be close to one, as suggested by the solution of the atmospheric neutrino problem (see Eqs.(24) and (25)), the lower bound $|U_{\mu 3}|_{(\min)}^2 = \frac{1}{2} \left(1 - \sqrt{1 - \sin^2 2\vartheta_{\mu\tau}^{(\min)}} \right)$ is close to 1/2.

If matter effects are important, the extraction of an upper bound for $|U_{e3}|^2$ from the data of $\nu_\mu \rightarrow \nu_e$ accelerator long-baseline experiments is more complicated. In this case the probability of $\nu_\mu \rightarrow \nu_e$ oscillations is given by (see [24])

$$P_{\nu_\mu \rightarrow \nu_e} = \frac{4|U_{e3}|^2 |U_{\mu 3}|^2}{\left(1 - \frac{A_{CC}}{\Delta m_{31}^2}\right)^2 + 4|U_{e3}|^2 \frac{A_{CC}}{\Delta m_{31}^2}} \sin^2 \left(\frac{\Delta m_{31}^2 L}{4E} \sqrt{\left(1 - \frac{A_{CC}}{\Delta m_{31}^2}\right)^2 + 4|U_{e3}|^2 \frac{A_{CC}}{\Delta m_{31}^2}} \right), \quad (37)$$

where E is the neutrino energy and L is the distance of propagation. This probability depends on the neutrino energy not only through the explicit E in the denominator of the phase, but also through the energy dependence of $A_{CC} \equiv 2EV_{CC}$. For long-baseline neutrino beams propagating in the mantle of the Earth the charged-current effective potential $V_{CC} = \sqrt{2}G_F N_e$ is practically constant: $N_e \simeq 2 N_A \text{ cm}^{-3}$ (N_A is the Avogadro number) and $V_{CC} \simeq 1.5 \times 10^{-13} \text{ eV}$.

If long-baseline experiments will not observe $\nu_\mu \rightarrow \nu_e$ transitions (or will find that they have an extremely small probability) for neutrino energies such that $A_{CC} \lesssim \Delta m_{31}^2$, it will

mean that $|U_{e3}|^2$ is small and a fit of the experimental data with the formula (37) will yield a stringent upper limit for $|U_{e3}|^2$ (taking into account the lower limit $|U_{\mu 3}|^2 \geq |U_{\mu 3}|_{(\min)}^2$ obtained from the solution of the atmospheric neutrino anomaly and from the observation of $\nu_\mu \rightarrow \nu_\tau$ long-baseline transitions). On the other hand, the non-observation of $\nu_\mu \rightarrow \nu_e$ transitions for neutrino energies such that $A_{CC} \gg \Delta m_{31}^2$ does not provide any information on $|U_{e3}|^2$ because in this case the transition probability (37) is suppressed by the matter effect. Hence, in order to check the hypothesis $|U_{e3}| \ll 1$, as well as to have some possibility to observe $\nu_\mu \rightarrow \nu_e$ transitions if this hypothesis is wrong, it is necessary that a substantial part of the energy spectrum of the neutrino beam lies below

$$\frac{\Delta m_{31}^2}{2V_{CC}} \simeq 30 \text{ GeV} \left(\frac{\Delta m_{31}^2}{10^{-2} \text{ eV}^2} \right). \quad (38)$$

This requirement will be satisfied in the accelerator long-baseline experiments under preparation (K2K [40], MINOS [41], ICARUS [42] and others [43]) if Δm_{31}^2 is not much smaller than 10^{-2} eV^2 .

VI. CONCLUSIONS

We have considered the scheme of mixing of three neutrinos with the mass hierarchy (1) and with Δm_{21}^2 and Δm_{31}^2 relevant, respectively, for the oscillations of solar and atmospheric neutrinos.

We have shown that in the framework of this scheme the recent results of the CHOOZ experiment [25] imply that $|U_{e3}|^2$ is small, the oscillations of solar neutrinos are described by the two-generation formalism and the oscillations of solar and atmospheric neutrinos depend on different and independent elements of the neutrino mixing matrix, *i.e.* they decouple. We have also shown that if not only $|U_{e3}|^2 \ll 1$ but also $|U_{e3}| \ll 1$, then the oscillations of atmospheric neutrinos do not depend on matter effects and are also described by the two-generation formalism. In this case, with the identifications (10) and (23) the two-generation analyses of solar and atmospheric neutrino data provide direct information on the mixing parameters of three neutrinos (see Eqs.(30)–(32)).

Let us notice that the independence of the oscillations of atmospheric neutrinos from matter effects can be checked by comparison of the transition probabilities of neutrinos and antineutrinos in future accelerator long-baseline experiments.

If future results from reactor (CHOOZ [25], Palo Verde [38], Kam-Land [39]) and accelerator (K2K [40], MINOS [41], ICARUS [42] and others [43]) long-baseline neutrino oscillation experiments will confirm and improve the upper bound (8) for $|U_{e3}|^2$ obtained from the first results of the CHOOZ experiment [25], the indications in favor of a decoupling of solar and atmospheric neutrino oscillations and of their accurate description by the two-generation formalism will be strengthened. In this case the distinction of the three allowed solutions of the solar neutrino problem (small and large mixing angle MSW and vacuum oscillations), which is a goal of the new generation of solar neutrino experiments (SuperKamiokande, SNO, ICARUS, Borexino, GNO and others [37]) could provide an indication of the actual values of the elements of the neutrino mixing matrix selecting one of the three possibilities (30)–(32).

ACKNOWLEDGMENTS

C.G. would like to thank V. Berezhinsky for a stimulating discussion on neutrino physics and astrophysics.

REFERENCES

- [1] See: V. Berezhinsky, Invited lecture at the 25th *International Cosmic Ray Conference*, Durban, 28 July – 8 August, 1997, preprint astro-ph/9710126.
- [2] See: T. Gaisser, Proc. of *Neutrino 96*, Helsinki, June 1996, edited by K. Enqvist *et al.*, p. 211 (World Scientific, Singapore, 1997).
- [3] K. Inoue, Talk presented at the 5th International Workshop on *Topics in Astroparticle and Underground Physics*, Gran Sasso, Italy, September 1997 (http://www-sk.icrr.u-tokyo.ac.jp/doc/sk/pub/pub_sk.html); R. Svoboda, Talk presented at the Conference on *Solar Neutrinos: News About SNU's*, 2–6 December 1997, Santa Barbara, California (<http://www.itp.ucsb.edu/online/snu/>).
- [4] Y. Suzuki, Talk presented at the *International School of Nuclear Physics, Neutrinos in Astro, Particle and Nuclear Physics*, Erice, Italy, September 1997 (http://www-sk.icrr.u-tokyo.ac.jp/doc/sk/pub/pub_sk.html); E. Kearns, Talk presented at the Conference on *Solar Neutrinos: News About SNU's*, 2–6 December 1997, Santa Barbara, California (<http://www.itp.ucsb.edu/online/snu/>).
- [5] B.T. Cleveland *et al.*, Nucl. Phys. B (Proc. Suppl.) **38**, 47 (1995).
- [6] K.S. Hirata *et al.*, Phys. Rev. D **44**, 2241 (1991).
- [7] GALLEX Coll., W. Hampel *et al.*, Phys. Lett. B **388**, 384 (1996).
- [8] J.N. Abdurashitov *et al.*, Phys. Rev. Lett. **77**, 4708 (1996).
- [9] Y. Fukuda *et al.*, Phys. Lett. B **335**, 237 (1994).
- [10] R. Becker-Szendy *et al.*, Nucl. Phys. B (Proc. Suppl.) **38**, 331 (1995).
- [11] W.W.M. Allison *et al.*, Phys. Lett. B **391**, 491 (1997).
- [12] S.M. Bilenky and B. Pontecorvo, Phys. Rep. **41**, 225 (1978).
- [13] S.M. Bilenky and S.T. Petcov, Rev. Mod. Phys. **59**, 671 (1987).
- [14] C.W. Kim and A. Pevsner, *Neutrinos in Physics and Astrophysics*, Contemporary Concepts in Physics, Vol.8 (Harwood Academic Press, Chur, Switzerland, 1993).
- [15] N. Hata and P. Langacker, Phys. Rev. D **56**, 6107 (1997).
- [16] G.L. Fogli, E. Lisi and D. Montanino, preprint hep-ph/9709473.
- [17] M.C. Gonzalez-Garcia *et al.*, preprint hep-ph/9801368.
- [18] M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, ed. F. van Nieuwenhuizen and D. Freedman (North Holland, Amsterdam, 1979), p.315; T. Yanagida, Proc. of the *Workshop on Unified Theory and the Baryon Number of the Universe*, KEK, Japan, 1979; R.N. Mohapatra and G. Senjanović, Phys. Rev. Lett. **44**, 912 (1980).
- [19] X. Shi and D.N. Schramm, Phys. Lett. B **283**, 305 (1992).
- [20] G.L. Fogli, E. Lisi and D. Montanino, Phys. Rev. D **49**, 3626 (1994); Astrop. Phys. **4**, 177 (1995); Phys. Rev. D **54**, 2048 (1996); G.L. Fogli, E. Lisi, D. Montanino and G. Scioscia, Phys. Rev. D **55**, 4385 (1997).
- [21] S. Goswami, K. Kar and A. Raychaudhuri, Int. J. Mod. Phys. A **12**, 781 (1997).
- [22] M. Narayan, M.V.N. Murthy, G. Rajasekaran and S.U. Sankar, Phys. Rev. D **53**, 2809 (1996); M. Narayan, G. Rajasekaran and S.U. Sankar, Phys. Rev. D **56**, 437 (1997); preprint hep-ph/9712409.
- [23] S.M. Bilenky, C. Giunti and C.W. Kim, Astrop. Phys. **4**, 241 (1996).
- [24] C. Giunti, C.W. Kim and M. Monteno, preprint hep-ph/9709439, to be published in Nucl. Phys. B.

- [25] M. Apollonio *et al.*, preprint hep-ex/9711002.
- [26] C. Athanassopoulos *et al.*, Phys. Rev. Lett. **77**, 3082 (1996).
- [27] S.M. Bilenky, C. Giunti and W. Grimus, Proc. of *Neutrino 96*, Helsinki, June 1996, edited by K. Enqvist *et al.*, p. 174 (World Scientific, Singapore, 1997); Eur. Phys. J. C **1**, 247 (1998).
- [28] N. Okada and O. Yasuda, Int. J. Mod. Phys. A **12**, 3669 (1997).
- [29] S.M. Bilenky, C. Giunti, C.W. Kim and M. Monteno, preprint hep-ph/9711400.
- [30] J.N. Bahcall and M.H. Pinsonneault, Rev. Mod. Phys. **67**, 781 (1995);
- [31] S. Turck-Chièze *et al.*, Phys. Rep. **230**, 57 (1993).
- [32] V. Castellani *et al.*, Phys. Rep. **281**, 309 (1997).
- [33] S.P. Mikheyev and A.Yu. Smirnov, Yad. Fiz. **42**, 1441 (1985) [Sov. J. Nucl. Phys. **42**, 913 (1985)]; Il Nuovo Cimento C **9**, 17 (1986).
- [34] T.K. Kuo and J. Pantaleone, Rev. Mod. Phys. **61**, 937 (1989).
- [35] G.L. Fogli, E. Lisi, A. Marrone and D. Montanino, preprint hep-ph/9711421.
- [36] L. Maiani, Proc. of the *International Symposium on Lepton Interactions at High Energies*, (Desy, Hamburg, 1977) p.867; see also: Particle Data Group, R.M. Barnett *et al.*, Phys. Rev. D **54**, 1 (1996).
- [37] Y. Suzuki (SuperKamiokande), Proc. of the *Fourth International Solar Neutrino Conference*, Heidelberg, Germany, 8–11 April 1997, edited by W. Hampel (Max-Planck-Institut für Kernphysik, 1997), p.163; R. Meijer Drees (SNO), *ibid.*, p.210; F. von Feilitzsch (Borexino), *ibid.*, p.192; E. Bellotti (GNO), *ibid.*, p.173; K. Lande (Homestake Iodine), *ibid.*, p.228; C. Tao (HELLAZ), *ibid.*, p.238; A.V. Kopylov (Lithium), *ibid.*, p.263; Yu.G. Zdesenko (Xenon), *ibid.*, p.283; P. Cennini *et al.* (ICARUS), LNGS-94/99-I, May 1994; T.J. Bowels (GaAs), Proc. of *Neutrino 96*, Helsinki, Finland, 13–19 June 1996, edited by K. Enqvist, K. Huitu and J. Maalampi (World Scientific, Singapore, 1997), p.83.
- [38] F. Boehm *et al.*, *The Palo Verde experiment*, 1996 (<http://www.cco.caltech.edu/~songhoon/Palo-Verde/Palo-Verde.html>).
- [39] F. Suekane, preprint TOHOKU-HEP-97-02, June 1997.
- [40] Y. Suzuki, Proc. of *Neutrino 96*, Helsinki, June 1996, edited by K. Enqvist *et al.*, p. 237 (World Scientific, Singapore, 1997).
- [41] MINOS Coll., D. Ayres *et al.*, NUMI-L-63, February 1995.
- [42] ICARUS Coll., P. Cennini *et al.*, LNGS-94/99-I, May 1994.
- [43] NOE Coll., G.C. Barbarino *et al.*, INFN/AE-96/11, May 1996; AQUA-RICH Coll., T. Ypsilantis *et al.*, LPC/96-01, CERN-LAA/96-13; OPERA Coll., H. Shibuya *et al.*, CERN-SPSC-97-24.